arXiv:hep-th/0509115v1 15 Sep 2005

ON THE POSSIBILITY OF QUANTUM-MECHANICAL INTERPRETATION THE RELATIVISTIC EFFECT OF ENERGY INCREASE IN A PARTICLE FREELY MOVING IN VACUUM

G. N. Gestrina
Usikov Institute of Radio Physics and Electronics
of National Academy of Ukraine
12, Proskura st., Kharkov, 61085, Ukraine
E-mail: ire@ire.kharkov.ua

Abstract

The relativistic effect of energy increase in a particle freely moving in vacuum is discussed on the basis of quantum field theory and probability theory using some ideas of super-symmetrical theories. The particle is assumed to consist of a "seed" whose energy is equal to the particle rest energy and whose pulse is equal to the product of the particle mass by its velocity and of a "fur coat" - the system of virtual quanta of the material field - vacuum. Each of these quanta possesses the same energy and pulse as the "seed" but have no mass. The system of the quanta is in a state being the superposition of quantum states with energies and pulses multiple of the "seed" energy and pulse. The virtual quanta is created (or destroyed) in of such states. The probability of creating a quanta in any state is the inverse of the relativistic factor, and the average number of the quanta making up the "fur coat" with a "seed" is equal to this particular factor. The kinetic energy and the relativistic addition to the particle pulse are interpreted as the average magnitude of the energy and the pulse in the system of the virtual quanta that constitute the particle "fur coat".

The Einstein formula linking the energy and the mass of a particle freely moving in vacuum, certainly, is the central formula of special theory of relativity [1]. It has played a tremendous role for a deeper understanding of the material world surrounding us and has had an enormous practical application. This formula shows that the energy of a relativistic particle increases with its velocity whereas its mass remains constant [2].

On the modern stage of physical science it becomes possible to describe this effect by means the quantum field theory. In this case some ideas of super-symmetrical theories turned out to be rather useful [3].

1. First it will be shown how by means of a simple mathematical transformation the Einstein formula may be expressed in the form allowing its interpretation on the quantum-mechanical basis.

The kinetic energy of a relativistic particle with mass m moving with constant velocity $\mathbf{v}$ may be presented as follows [4]:

$$E_{kin} = mc^2\left(\frac{1}{\sqrt{1-\left(\frac{v}{c}\right)^2}} - 1\right) = = mc^2\sqrt{1-\left(\frac{v}{c}\right)^2}\,\frac{1-\sqrt{1-\left(\frac{v}{c}\right)^2}}{1-\left(\frac{v}{c}\right)^2} = = mc^2\sqrt{1-\left(\frac{v}{c}\right)^2}\sum_{n=1}^{\infty} n\left[1-\sqrt{1-\left(\frac{v}{c}\right)^2}\right]^n , \tag{1}$$

[1]

where c is velocity of light in vacuum. Supposing further that

$$E_n = nmc^2 \quad (n = 1, 2, \ldots) \tag{2}$$

and

$$P_n = P_0(1-P_0)^n \tag{3}$$

$$(n = 0, 1, 2, \ldots,\ 0 < P_n < 1,\ \sum_{n=0}^{\infty} P_n = 1),$$

where

$$P_0 = \sqrt{1-\left(\frac{v}{c}\right)^2}, \tag{4}$$

we get

$$E_{kin} = \sum_{n=1}^{\infty} E_n P_n, \tag{5}$$

The full energy E of a relativistic particle can be written as:

$$E = mc^2 + \sum_{n=1}^{\infty} E_n P_n, \tag{6}$$

and its pulse $\mathbf{p}$ due to the relativistic relationship $\mathbf{p} = E\mathbf{v}/c^2$ will take the form:

$$\mathbf{p} = m\mathbf{v} + \sum_{n=1}^{\infty} \mathbf{p}_n P_n, \tag{7}$$

where

$$\mathbf{p}_n = nm\mathbf{v} \quad (n = 1, 2, \ldots). \tag{8}$$

As it follows from relations (6) and (7) a relativistic particle with mass m freely moving in vacuum at constant velocity $\mathbf{v}$, can be considered as an aggregate consisting of a "seed" particle with energy $E_0 = mc^2$ and pulse $\mathbf{p_0} = m\mathbf{v}$ and an infinite system of virtual quanta of the material field, i.e., the vacuum ("fur coat"), each of them also having energy $E_0 = mc^2$ and pulse $\mathbf{p}_0 = m\mathbf{v}$. This system of quanta is in a state representing the superposition of n^{thes} quantum states, characterized by energies $E_n = nmc^2$ and pulses $\mathbf{p}_n = nm\mathbf{v} \quad (n = 1, 2, \ldots)$.

The interpretation of P_n values as some probabilities will be given below. The accumulation by the "seed" particle of such a "fur coat" consisting of virtual quanta is very much alike the crystallization process in a saturated solution of a substance after an isolated crystal "seed" of the same substance has been introduced into it. In both cases a similar reaction of the ambient medium (a vacuum or a saturated solution) is observed with the only difference that in the first case virtual field quanta possessing the same energy and pulses have no mass whereas crystals obtained from the saturated solution represent precise copies of "seed", i.e., of an isolated crystal introduction from outside.

Consequently, already within the framework of the special relativity theory vacuum acquires the sense of a material field. The virtual quanta of this field are bosons and n^{thes} quantum states of the virtual quantum quantum system are n^{thes} boson states.

2. Let us correlate each n^{th} boson state of this system with the n^{th} excited state of a harmonic oscillator with frequency $\omega = E_0/\hbar$ ($\hbar = h/2\pi$, h–Plank constant). However in the situation described the lowest energy state of the material field for a particle (i.e., the vacuum) has a zero energy whereas the lowest energy value of the harmonic oscillator is $\hbar\omega/2$. To obtain a correspondence between the bosonic states of virtual quanta and the excited state of the harmonic oscillator through rejecting $\hbar\omega/2$ energy is impossible to neglect zero oscillations of the vacuum since they are wholly real [3].

In super-symmetrical theories where bosons and fermions are considered together the infinite positive energy of zero boson oscillations is fully compensated by the infinite negative energy of fermion oscillations therefore the energy of a single boson-fermion vacuum is zero. What is considered in super-symmetrical theories is a super-symmetrical harmonic oscillator. Hamiltonian of such an oscillator is the sum of Hamiltonians of two non-interacting harmonic oscillators corresponding to boson and fermion degrees of freedom. Frequencies of these oscillators must be equal. The lowest energetic state of the super-symmetrical oscillator is zero [3].

In some situations when fermion degrees of freedom are absent there exists, however, the possibility to use the ideas concerning the zero energy of the vacuum and two harmonic oscillators with equal frequencies. Thus, in the described situation it is sufficient to suppose that in the n^{th} boson state there are $n_+ = N_n + n$ bosons with $E_0 > 0$ and $n_- = N_n$ bosons with $E_0 < 0$ ($E_0 = \pm mc^2$). The difference n ($n = n_+ - n_-$) determines the energy of the n^{th} boson state. N_n virtual quanta with $E_0 > 0$ compensate N_n ones with $E_0 < 0$. N_n is a random positive integer therefore all the n^{ths} quantum boson states are degenerate.

Let the basis vector of the n^{ths} quantum states of the virtual quantum system be represented in the form [3]:

$$|n_+, n_-\rangle = |\psi_n\rangle. \tag{9}$$

Here n_+ and n_- are population numbers for quanta with $E_0 > 0$ and $E_0 < 0$, respectively, $|\psi_n\rangle$ is the vector of the n^{th} quantum state. $|N_n, N_n\rangle$ vector will describe the vacuum field state. It is quite clear that the energy of the vacuum is zero.

Now it is necessary to consider two harmonic oscillators with the same frequency $\omega = |E_0|/\hbar$. Let q_1, p_1', m_1 and H_+ be the generalized coordinate, the pulse, the mass and the Hamiltonian of the first oscillator corresponding to boson degrees of freedom with $E_0 > 0$, and q_2, p_2', m_2 and H_- be the generalized coordinate, the pulse, the mass and the Hamiltonian of the second oscillator corresponding to boson degrees of freedom with $E_0 < 0$, that is

$$H_+ = \frac{1}{2}\left(\frac{p_1^{'2}}{m_1^2} + m_1 \omega q_1^2\right), \; m_1 = +\frac{|E_0|}{c^2} \tag{10}$$

and

$$H_- = \frac{1}{2}\left(\frac{p_2^{'2}}{m_2^2} + m_2 \omega q_2^2\right), \; m_2 = -\frac{|E_0|}{c^2}. \tag{11}$$

Correspondingly the energy levels of these oscillators have the form

$$E_{n_+} = \hbar\omega(n_+ + \frac{1}{2}) \quad (n_+ = n, n+1, \ldots) \tag{12}$$

and

$$E_{n_-} = \hbar\omega(n_- + \frac{1}{2}) \quad (n_- = 0, 1, \ldots) \tag{13}$$

Creation and destruction operators for bosons with $E_0 > 0 \;\; (C^{\pm}_{+})$ and for bosons with $E_0 < 0 \;\; (C^{\pm}_{-})$ are defined as follows [5]:

$$C^{\pm}_{+} = (2\hbar\omega m_1)^{-1/2}(p'_1 \pm i m_1 \omega q_1) \tag{14}$$

and

$$C^{\pm}_{-} = (-2\hbar\omega m_2)^{-1/2}(p'_2 \pm i m_2 \omega q_2). \tag{15}$$

They satisfy the ordinary commutation relations

$$[C^{-}_{\pm}, C^{+}_{\pm}] = 1, \quad [C^{\pm}_{-}, C^{\pm}_{+}] = 0. \tag{16}$$

Their effect on state vectors $|n_+, n_-\rangle$ will be the following:

$$C^{+}_{+}|n_+, n_-\rangle = \sqrt{n_+ + 1}\;|n_+ + 1, n_-\rangle; \tag{17}$$

$$C^{-}_{+}|n_+, n_-\rangle = \sqrt{n_+}\;|n_+ - 1, n_-\rangle; \tag{18}$$

$$C^{+}_{-}|n_+, n_-\rangle = \sqrt{n_+ 1}\;|n_+, n_- + 1\rangle; \tag{19}$$

$$C^{-}_{-}|n_+, n_-\rangle = \sqrt{n_-}\;|n_+, n_- - 1\rangle. \tag{20}$$

Thus, the creation operator C^{+}_{+} acting on $|n_+, n_-\rangle = |\psi_n\rangle$ vector transforms it to $|n_+ + 1, n_-\rangle = |\psi_{n+1}\rangle$ vector, whereas the annihilation operator C^{-}_{+} transforms the same vector to $|n_+ - 1, n_-\rangle = |\psi_{n-1}\rangle$ vector.

Similarly the creation operator C^{+}_{-} acting on $|n_+, n_-\rangle = |\psi_n\rangle$ vector transforms it into $|n_+, n_- + 1\rangle = |\psi_{n-1}\rangle$ vector, at the same time the annihilation C^{-}_{-} operator transforms the same vector into $|n_+, n_- - 1\rangle = |\psi_{n+1}\rangle$ vector.

It can be seen that $|n_+ + 1, n_-\rangle$ and $|n_+, n_- - 1\rangle$ vectors describe equivalent energy states characterized by $|\psi_{n+1}\rangle$ vector, and $|n_+ - 1, n_-\rangle$ and $|n_+, n_- + 1\rangle$ vectors describe those characterized by $|\psi_{n-1}\rangle$ vector.

Operator

$$C^{+}_{+}C^{-}_{+} = \hat{n}_+ \tag{21}$$

will be an operator of the boson number with $E_0 > 0$, whereas operator

$$C^{+}_{-}C^{-}_{-} = \hat{n}_- \tag{22}$$

will define the number of bosons with $E_0 < 0$.

Correspondingly $H_\pm$ Hamiltonians will be expressed through $\hat{n}_\pm$ operators as follows:

$$H_+ = \hbar\omega(\hat{n}_+ + \frac{1}{2}) \tag{23}$$

and

$$H_- = -\hbar\omega(\hat{n}_- + \frac{1}{2}). \tag{24}$$

Since there is no interaction between the oscillators their common Hamiltonian H will be equal to the sum of H_+ and H_- Hamiltonians, i.e.,

$$H = \hbar\omega(\hat{n}_+ - \hat{n}_-). \tag{25}$$

The oscillator corresponding to this Hamiltonian is not super-symmetrical (in a super-symmetrical harmonic oscillator all the states except zero one are doubly degenerate [3]).

It is obvious that

$$H\;|\psi_n\rangle = E_n\;|\psi_n\rangle \quad (n = 1, 2, \ldots). \tag{26}$$

This means that each of $|\psi_n\rangle$ vectors is an eigenvector of H Hamiltonian pertaining to the eigenvalue of $E_n = nE_0$ energy.

The $|\psi_n\rangle$ vectors form an orthonormal system of vectors:

$$\langle\psi_n|\psi_{n'}\rangle = \delta_{nn'} \tag{27}$$

($\delta_{nn'}$ is the Kroneker symbol). Any admissible state $|\psi\rangle$ vector of virtual quantum system is a superposition of $|\psi_n\rangle$ vectors, i.e.,

$$|\psi\rangle = \sum_n \alpha_n\;|\psi_n\rangle, \tag{28}$$

where α_n factors are determined from relations

$$\alpha_n = \langle\psi|\psi_n\rangle \tag{29}$$

and satisfy the normalizing condition:

$$\sum_n |\alpha_n|^2 = 1. \tag{30}$$

The average value of virtual quantum energy in the state described by $|\psi\rangle$ vector equal:

$$\bar{E} = \langle\psi|H|\psi\rangle = \sum_n |\alpha_n|^2 E_n. \tag{31}$$

3. Let us clear up the meaning of P_n-values and their connection with α_n. For this purpose a complete system of independent events A_n and their opposite events $\bar{A}_n$

will be introduced (Bernoulli's scheme)[6]. A_n-event will mean the creation of a virtual quantum with $E_0 > 0$ (or the destruction of a virtual quantum with $E_0 < 0$) in the n^{th} quantum state with P_0 probability. At the same time let us assume that $\bar{A}_n$ means the destruction of a virtual quantum with $E_0 > 0$ (or the creation of a virtual quantum with $E_0 < 0$) in the n^{th} quantum state with a probability of $1 - P_0$.

$P_n = P_0(1 - P_n)^n$ values will mean the probabilities of that the creation of the virtual quantum with $E_0 > 0$ has not taken place in quantum states from 0 to $n - 1$ but only in the n^{th} quantum state (or the probability that the quantum with $E_0 < 0$ was not destructed in quantum states from 0 to $(n-1)^{th}$ but was destructed only in the n^{th} quantum state).

Let us determine the value

$$\bar{n}+1 = \sum_{n=1}^{\infty} nP_n + 1 = P_0 \sum_{n=1}^{\infty} n(1-P_0)^{n-1} = 1/P_0. \quad (32)$$

If $\bar{n}+1$ is multiplied by mc^2 one gets the complete energy E of a relativistic particle, whereas the production of the multiplying $\bar{n}+1$ by $m\mathbf{v}$ gives its pulse. Multiplying $\bar{n}$ by mc^2 we get an expression for E_{kin} of the relativistic particle. Consequently $\bar{n}$ value can be interpreted as an average number of virtual quanta forming "fur coat" of the relativistic particle whereas E_{kin} is an average value of energy of these quanta. Comparing the expression for E_{kin} (5) with expression (31) for the average value of energy of virtual quanta in a state that is a superposition of the n^{ths} quantum states described by $|\psi\rangle$ vector one can conclude that

$$|\alpha_n| = \sqrt{P_n}. \quad (33)$$

4. Let us write the expression for the de'Broglie wave of the relativistic particle as follows:

$$\begin{aligned} \psi &= C\exp[-i(Et - \mathbf{pr}/\hbar)] = \quad (34) \\ &C\exp\{-i[(E_0t - \mathbf{p}_0\mathbf{r}) + \sum_{n=1}^{\infty}(E_nt - \mathbf{p}_n\mathbf{r})P_n]/\hbar\} = \\ &= \psi_0 \prod_{n=1}^{\infty} \psi_n, \end{aligned}$$

where

$$\psi_0 = C_0\exp[-i(E_0t - \mathbf{p}_0\mathbf{r})/\hbar] \quad (35)$$

is the de'Broglie wave of the "seed" particle and

$$\begin{aligned} \psi_n &= C_n\exp[-i(E_nt - \mathbf{p}_n\mathbf{r})/\hbar] = \quad (36) \\ &= C_n\exp[-i(E_0t - \mathbf{p}_0\mathbf{r})nP_n/\hbar] \end{aligned}$$

are the de'Broglie waves of virtual quanta,

$$C = C_0 \prod_{n=1}^{\infty} C_n, \quad (37)$$

$\mathbf{r}$ is a radius-vector of any point of the space.

Expression (34) shows that there is no interaction between the "seed" and the virtual quanta as well as between the latters themselves.

All the de'Broglie waves ψ_0 and ψ_n $(n = 1, 2, \ldots)$ have the same phase velocity $v_p = c^2/v$, the frequencies $\omega_0 = E_0/\hbar$ are $\omega_n = E_nP_n/\hbar = n\omega_0P_n$ $(n = 1, 2, \ldots)$ and the lengths $\lambda_0 = \hbar/p_0$ and $\lambda_n = \hbar/p_nP_n$ $(n = 1, 2, \ldots)$, where ω_0 and λ_0 are the frequency and the de'Broglie wavelength of the "seed" particle whereas p_0 is its pulse.

But neither the "seed" particle nor the virtual quanta can be separately fixed by a device. The latter fixes the relativistic particle as a whole. This follows from the fact that diameters of the diffraction coils on the electronogram are expressed through the relativistic value for the de'Broglie wavelength $\lambda = \hbar\sqrt{1 - (\frac{v}{c})^2}/mv$ of a particle.

Thus the quantum field theory combined with some ideas of modern super-symmetrical theories allows to consider "fur coat" of a relativistic particle freely moving in vacuum as a virtual quantum collection of the vacuum each of which has the same energy and pulse as the "seed" particle.

The boson degrees of freedom discussed above are probably those internal degrees of freedom which P. Dirak [7, 8] had in mind suggesting the availability of the dynamic structure in the relativistic particle.

From the standpoint of the probability theory the "fur coats" formation of the relativistic particle out of vacuum virtual quantum is regarded as a random event.

1. A. Einstein, Ann. d. Phys. Bd. 17, 851 (1905).
2. L.B. Okun', UFN, 158, 511 (1989).
3. L.E. Gendenshtein, I.V. Krive, UFN, 146, 553 (1985)
4. G.N. Gestrina, Reports of Uk AS, ser "A", 9, 55 (1990)
5. V.B.Berestetsky, E.M.Lifshits, L.P.Pitaevsky. Quantum Electrodynamics. Moscow; Nauka, 1974 (in Russian)
6. E.S.Ventsel. Theory Probabilities. Moscow; Nauka, 1984 (in Russian)
7. P.A.M. Dirac, Proc.Roy.Soc., A,322,435 (1971)
8. P.A.M. Dirac, Proc.Roy.Soc., A,328,1 (1972)